

IMAGES IN...

Cranial thoracic stenotic spondylomyelopathy in a young Rottweiler

Silke Hecht,¹ Jennifer Michaels,^{1,2} Sarah Childers,¹ Linden Craig³

¹Small Animal Clinical Sciences, University of Tennessee, Knoxville, Tennessee, USA

²Angell Animal Medical Center, Boston, Massachusetts, USA

³Biomedical and Diagnostic Sciences, University of Tennessee, Knoxville, Tennessee, USA

Correspondence to

Silke Hecht; shecht@utk.edu

Received 9 September 2016

Revised 11 January 2017

Accepted 16 January 2017

DESCRIPTION

A four-month-old male entire Rottweiler presented for progressive pelvic limb ataxia. Neurological examination revealed non-ambulatory paraparesis with ataxia in pelvic limbs, absent pelvic limb postural reactions, clonic patellar reflexes, normal withdrawal reflexes in all limbs and hyperaesthesia on neck manipulation. Weight shifting and splaying of the thoracic limbs (representing either mild paresis or compensatory gait change) were also noted. Neurolocalisation was third thoracic to third lumbar spinal cord segments (Th3-L3), most likely within the cranial aspect of this region. MRI showed enlargement of the dorsal laminae and articular processes of Th1-4, causing multifocal vertebral canal stenosis and attenuation of epidural fat and cerebrospinal fluid (CSF) signal on T2-weighted and myelographic sequences (Fig 1a,b). Changes were most severe at Th2-3 and associated with

moderate dorsoventral spinal cord compression (Fig 1c). Lesser degree spinal cord compression was noted at Th1-2 and Th3-4 (Fig 1d,e). CT confirmed the MRI findings (Fig 1f). The imaging diagnoses were cranial thoracic stenosis due to vertebral malformation and articular process/dorsal lamina hypertrophy previously reported in large and giant breed dogs (Stalin and others 2009, Johnson and others 2012). Surgical decompression (dorsal laminectomy) was declined. Postmortem evaluation yielded a final diagnosis of developmental thoracic stenotic spondylomyelopathy with secondary severe segmental chronic myelomalacia (Fig 2). Cranial thoracic stenosis most commonly affects young, heavy, male dogs with a conformation typical of Molosser breeds. This disorder is less common than osseous-associated cervical spondylomyelopathy but is similar with regard to patient signalment and morphological features (da Costa 2010).




FIG 1: This four-month-old male Rottweiler presented with progressive ataxia. MRI and CT were performed. MR images of the cervicothoracic spine. Sagittal T2-weighted (a) and myelographic (b) images showing vertebral canal stenosis and attenuation of dorsal epidural fat and cerebrospinal fluid signal at Th2-3 and to a lesser degree Th1-2 and Th3-4 (arrows). Transverse T2-weighted image at Th2-3 (c) demonstrates enlargement of the articular processes as well as the dorsal lamina with marked narrowing of the vertebral canal (arrowheads), complete circumferential attenuation of epidural and subarachnoid space signal and moderate dorsoventral spinal cord compression. Lesser degree spinal cord compression due to articular process hypertrophy is also noted at Th1-2 and Th3-4 (d,e; arrowheads). (f) 3D rendering of CT images shows multifocal narrowing of the vertebral canal from dorsolaterally.



To cite: Hecht S, Michaels J, Childers S, et al. *Vet Rec Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/vetreccr-2016-000382




FIG 2: This is the cranial end of the third thoracic vertebra. The left cranial facet is more ventrally angled than the right. There is marked synovial and fibrous proliferation dorsal to the spinal canal resulting in a narrowed canal.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Additional patient data can be made available to interested readers upon request.

© British Veterinary Association (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- da Costa R. C. (2010) Cervical spondylomyelopathy (wobbler syndrome) in dogs. *Veterinary Clinics of North America: Small Animal Practice* 40, 881–913
- Johnson P., De Risio L., Sparkes A., McConnell F., Holloway A. (2012) Clinical, morphologic, and morphometric features of cranial thoracic spinal stenosis in large and giant breed dogs. *Veterinary Radiology and Ultrasound* 53, 524–534
- Stalin C., Pratt J., Smith P. M., Jefferey N. (2009) Thoracic stenosis causing lateral compression of the spinal cord in two immature dogues de bordeaux. *Veterinary and Comparative Orthopaedics and Traumatology* 22, 59–62

Copyright 2017 British Veterinary Association. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

Veterinary Record Case Reports subscribers may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit vetrecordcasereports.bvapublications.com for more articles like this and to become a subscriber

Cranial thoracic stenotic spondylomyelopathy in a young Rottweiler

Silke Hecht, Jennifer Michaels, Sarah Childers and Linden Craig

Vet Rec Case Rep 2017 5:
doi: 10.1136/vetreccr-2016-000382

Updated information and services can be found at:
<http://vetrecordcasereports.bmj.com/content/5/1/e000382>

References

These include:

This article cites 3 articles, 0 of which you can access for free at:
<http://vetrecordcasereports.bmj.com/content/5/1/e000382#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>